

45_CONCORSO PUBBLICO, PER TITOLI ED ESAMI, PER LA COPERTURA A TEMPO DETERMINATO, DELLA DURATA DI CINQUE ANNI PER N. 1 POSTO DI RICERCATORE SANITARIO DA ASSEGNARE ALLA SC NEUROLOGIA 8 – DEMENZE E PATOLOGIE DEGENERATIVE DEL SISTEMA NERVOSO CENTRALE

PROVA 1

1. L'atrofia corticale posteriore (PCA), è una sindrome clinica che può rappresentare la manifestazione di diverse malattie neurodegenerative. Indichi le principali patologie neurodegenerative che possono sostenerla e descriva le principali caratteristiche cliniche, neuropsicologiche e neuroradiologiche di una di esse
2. La sigla "ICT" è l'acronimo di:
 - a) Internet Computer Transfer
 - b) Internet Carrier Television
 - c) Information and Communication Technologies
3. Leggere e tradurre il seguente testo

ABSTRACT

The Dementia with Lewy Bodies (DLB) Consortium has refined its recommendations about the clinical and pathologic diagnosis of DLB, updating the previous report, which has been in widespread use for the last decade. The revised DLB consensus criteria now distinguish clearly between clinical features and diagnostic biomarkers, and give guidance about optimal methods to establish and interpret these. Substantial new information has been incorporated about previously reported aspects of DLB, with increased diagnostic weighting given to REM sleep behavior disorder and ^{123}I -iodine-metaiodobenzylguanidine (MIBG) myocardial scintigraphy. The diagnostic role of other neuroimaging, electrophysiologic, and laboratory investigations is also described. Minor modifications to pathologic methods and criteria are recommended to take account of Alzheimer disease neuropathologic change, to add previously omitted Lewy-



free DS M
Giuseppe Maggini
Sorom

PROVA NON ESTRATTA

31/07/2026

Carlo Ugo Uboldi

45_CONCORSO PUBBLICO, PER TITOLI ED ESAMI, PER LA COPERTURA A TEMPO DETERMINATO, DELLA DURATA DI CINQUE ANNI PER N. 1 POSTO DI RICERCATORE SANITARIO DA ASSEGNARE ALLA SC NEUROLOGIA 8 – DEMENZE E PATOLOGIE DEGENERATIVE DEL SISTEMA NERVOSO CENTRALE

PROVA 2

1. Descriva la differenza tra Corticobasal Syndrome (CBS) e Corticobasal Degeneration (CBD), riportando i principali correlati neuropatologici e le possibili diagnosi eziologiche associate alla CBS
2. Per SPAM si intende:
 - a) l'invio non richiesto di messaggi di posta elettronica con fini commerciali o offensivi
 - b) un programma che permette di scrivere messaggi di posta elettronica
 - c) una truffa bancaria on line
3. Leggere e tradurre il seguente testo

Based on the recent literature and collective experience, an international consortium developed revised guidelines for the diagnosis of behavioural variant frontotemporal dementia. The validation process retrospectively reviewed clinical records and compared the sensitivity of proposed and earlier criteria in a multi-site sample of patients with pathologically verified frontotemporal lobar degeneration. According to the revised criteria, 'possible' behavioural variant frontotemporal dementia requires three of six clinically discriminating features (disinhibition, apathy/inertia, loss of sympathy/empathy, perseverative/compulsive behaviours, hyperorality and dysexecutive neuropsychological profile). 'Probable' behavioural variant frontotemporal dementia adds functional disability and characteristic neuroimaging, while behavioural variant frontotemporal dementia 'with definite frontotemporal lobar degeneration' requires histopathological confirmation or a pathogenic mutation. Sixteen brain banks contributed cases meeting histopathological criteria for frontotemporal lobar degeneration and a clinical diagnosis of behavioural variant frontotemporal dementia, Alzheimer's disease, dementia with Lewy bodies or vascular dementia at presentation. Cases with predominant primary progressive aphasia or extra-pyramidal syndromes were excluded. In these autopsy-



[Handwritten signature]
Francesca Maggiani

PROVA ESTRATTA

31/07/2016

Cavaliere Marco Mollà

45_CONCORSO PUBBLICO, PER TITOLI ED ESAMI, PER LA COPERTURA A TEMPO DETERMINATO, DELLA DURATA DI CINQUE ANNI PER N. 1 POSTO DI RICERCATORE SANITARIO DA ASSEGNARE ALLA SC NEUROLOGIA 8 – DEMENZE E PATOLOGIE DEGENERATIVE DEL SISTEMA NERVOSO CENTRALE

PROVA 3

1. Descriva il profilo neuropsicologico tipico delle patologie neurodegenerative a prevalente interessamento sottocorticale. Quali funzioni cognitive valuterebbe e quali prove neuropsicologiche utilizzerebbe per indagare il rallentamento cognitivo, le funzioni esecutive, l'attenzione, la memoria e gli aspetti visuo-spaziali?
2. Il tasto CAPS LOCK sulla tastiera serve per:
 - a) attivare una modalità di inserimento del testo in cui le lettere sono sempre maiuscole
 - b) attivare il tastierino numerico usare
 - c) i tasti funzione
3. Leggere e tradurre il seguente testo

ABSTRACT

This article provides a classification of primary progressive aphasia (PPA) and its 3 main variants to improve the uniformity of case reporting and the reliability of research results. Criteria for the 3 variants of PPA—nonfluent/agrammatic, semantic, and logopenic—were developed by an international group of PPA investigators who convened on 3 occasions to operationalize earlier published clinical descriptions for PPA subtypes. Patients are first diagnosed with PPA and are then divided into clinical variants based on specific speech and language features characteristic of each subtype. Classification can then be further specified as “imaging-supported” if the expected pattern of atrophy is found and “with definite pathology” if pathologic or genetic data are available. The working recommendations are presented in lists of features, and suggested assessment tasks are also provided. These recommendations have been widely agreed upon by a large group of experts and should be used to ensure consistency of PPA classification in



[Handwritten signature]
Giuseppe Magliani

PROVA NON ESTRATTA

31/07/2016

Ceschi Ubaldo